I° Congresso Nazionale Gruppo di Studio SIN
Neurologia di Genere :
*Un diverso modo di pensare
alle malattie del sistema nervoso*

Roma, 4 Maggio 2019

Genetica e medicina di genere

Barbara Garavaglia PhD

Fondazione I.R.C.C.S.
Istituto Neurologico Carlo Besta

Sistema Socio Sanitario

Regione
Lombardia

Sex, Gender, Genetics, and Health

Susan E. Short, PhD, Yang Claire Yang, PhD, and Tania M. Jenkins, MA

This article addresses 2 questions. First, to what extent are sex and gender incorporated into research on genetics and health? Second, how might social science understandings of sex and gender, and gender differences in health, become more integrated into scholarship in this area? We review articles on genetics and health published in selected peer-reviewed journals. Although sex is included frequently as a control or stratifying variable, few articles articulate a conceptual frame or methodological justification for conducting research in this way, and most are not motivated by sex or gender differences in health. Gender differences in health are persistent, unexplained, and shaped by multilevel social factors. Future scholarship on genetics and health needs to incorporate more systematic attention to sex and gender, gender as an environment, and the intertwining of social and biological variation over the life course. Such integration will advance understandings of gender differences in health, and may yield insight regarding the processes and circumstances that make genomic variation relevant for health and well-being. (*Am J Public Health*. 2013;103:S93–S101. doi:10.2105/AJPH.2013.301229)

^aAmerican Journal of Epidemiology (AJE) publishes 2 issues/month, for 12 months (24 issues/year).
^bAmerican Journal of Public Health (AJPH) publishes 1 issue/month, for 12 months (12 issues/year).
^cEpidemiology publishes 1 issue every 2 months (6 issues/year).
^dNature Genetics publishes 1 issue/month, for 12 months (12 issues/year).

FIGURE 1—Selection process of articles reviewed, 2007–2011.

TABLE 1—Summary of Incorporation of Sex or Gender in Research Articles on Genetics and Health

Variable	All Journals, ^a No. (Proportion)	Public Health Journals, ^b No. (Proportion)
Total number of articles	198	139
Single sex/gender analysis sample		
Total	35	32
Reproductive or sex-specific outcome ^c	12 (0.34)	10 (0.31)
Non-sex-specific health outcome	23 (0.66)	22 (0.69)
Multiple sex/gender analysis sample		
Total	163	107
Level of inclusion		
Sex/gender in title	2 (0.01)	0 (0.00)
Sex/gender in abstract	19 (0.12)	18 (0.13)
Type of inclusion		
Match on sex in sampling	20 (0.12)	14 (0.10)
Control/adjust for sex in analysis	93 (0.57)	62 (0.45)
Incorporate sex with twin design	16 (0.10)	15 (0.11)
Stratify analysis by sex	34 (0.21)	22 (0.16)
Interact sex with genetic variant	12 (0.07)	6 (0.04)
No mention of treatment of sex/gender in analysis	28 (0.17)	10 (0.07)

Note. All articles were published between 2007 and 2011. Ten studies did not clearly indicate which sex(es) or gender(s) they included and are presumed not to be single sex or gender studies.
^aAll Journals includes American Journal of Epidemiology, American Journal of Public Health, Epidemiology and Nature Genetics.
^bPublic Health Journals excludes Nature Genetics.
^cBreast cancer was coded as non-sex-specific health outcome.

Supplement 1, 2013, Vol 103, No. S1 | American Journal of Public Health

I FATTI:

Le donne vivono più a lungo degli uomini ma si ammalano di più
(paradosso donna)

La dimensione e la composizione corporea è diversa tra maschi e femmine

La farmacocinetica e la farmacodinamica è diversa tra uomini e donne
(le donne presentano più eventi avversi degli uomini al consumo dei farmaci)

Tab. IV. Differenze di genere nell'attività degli Isoenzimi del Citocromo P450. *Gender differences in Cytochrome P450 Isoenzyme activity.*

Isoenzima P450	Genere femminile	Età senile	Farmaco
CYP3A4	↑ Attività	↓ Attività	BDZ, nefazodone
CYP1A2	↓ Attività	No effetto	Fluoxamina
CYP2C19	↑ Attività	No effetto	Citalopram, clomipramina

D. MARAZZITI, A. BALDINI, M. CARLINI ET AL. GIORN ITAL PSICOPAT 2007; 13: 69-75

Sistema Socio Sanitario

I FATTI

Molte malattie sono determinate da differenze biologiche legate al sesso:

- **Donne a maggior rischio di malattie autoimmuni**
- **Donne a maggior incidenza di fratture**
- **Donne a maggior rischio di depressione e demenza (AD), uomini a maggior rischio di Parkinson**
- **Uomini a maggior rischio di malattie cardiovascolari in età più giovane, dopo la menopausa aumento del rischio nelle donne**
- **Uomini a maggior rischio di mortalità per cancro**

Sistema Socio Sanitario

Fondazione I.R.C.C.S.
Istituto Neurologico Carlo Besta

Regione
Lombardia

Regitz-Zagrosek V, EMBO Rep. 2012;13:596-603

Sistema Socio Sanitario

Cross-sex hormone administration changes pain in transsexual women and men.

Aloisi AM¹, Bachiocco V, Costantino A, Stefani R, Ceccarelli I, Bertaccini A, Meriggola MC.

⊕ Author information

Abstract

Chronic pain is gender-related, since there is a clear predominance of one sex with respect to the other in most pain syndromes. Gonadal hormones are known to affect the occurrence and incidence of pain. Transsexuals receive cross-sex hormones to develop and maintain somatic characteristics of the opposite sex: male to female transsexuals (MtF) are administered estrogens and anti-androgens, while female to male transsexuals (FtM) are administered androgens. Hence, these subjects represent a model to study the relationship between sex hormones and pain. Questionnaires dealing with sociodemographic data and pain (occurrence, frequency, duration, intensity, location and associated symptoms) were administered to both MtF and FtM transsexuals under hormone treatment for sex reassignment for at least 1 year. Forty-seven MtF and 26 FtM completed the questionnaires. Fourteen of the 47 MtF (29.8%) reported painful conditions, which in 11 subjects were not present before the beginning of hormone treatment. Pain consisted mainly of headaches and breast and musculoskeletal pain. Five subjects suffered from more than one pain condition. Sixteen of the 26 FtM (61.5%) reported pain. In 11 subjects, the pain was present before the beginning of hormone intake, and in 6 of them it improved after testosterone administration. These data suggest that marked changes in sex hormones affect the occurrence of pain in a high percentage of humans but not in all of them. Whether these effects are due to peripheral or central actions of sex steroids is unknown.

PMID: 17379410 DOI: [10.1016/j.pain.2007.02.006](https://doi.org/10.1016/j.pain.2007.02.006)

Le donne sono
a maggior
rischio di
dolore
cronico

I transessuali che ricevono una terapia ormonale per sviluppare e mantenere caratteristiche somatiche del sesso opposto possono risultare più inclini o più protetti in base al trattamento ormonale:

- **un terzo dei maschi, transessuali femminili**, sviluppano dolore cronico in concomitanza con il trattamento a base di estrogeni/anti-androgeni e ha mostrato una maggiore sensibilità al dolore secondo il calendario di terapia ormonale
- **la metà delle femmine, transessuali maschi**, trattata con testosterone ha registrato un significativo miglioramento del dolore cronico (per esempio, mal di testa) già presente prima dell'inizio del trattamento e ha migliorato i profili di sensibilità al dolore in seguito alla terapia ormonale.

Sistema Socio Sanitario

Fondazione I.R.C.C.S.
Istituto Neurologico Carlo Besta

Regione
Lombardia

Genoma Umano

Circa 3,2 miliardi di basi
20-25.000 geni

Junk DNA: sequenze regolatorie, splicing alternativo, miRNA, ????

Sequenze codificanti proteine

Sistema Socio Sanitario

Fondazione I.R.C.C.S.
Istituto Neurologico Carlo Besta

Regione
Lombardia

EPIGENETICA: regola l'espressione genica senza alterare la sequenza del DNA

Meccanismi epigenetici:

Modificazioni del DNA = metilazione della citosina da parte delle metiltrasferasi

Modificazione delle proteine = acetilazione, metilazione, ubiquitinazione, fosforilazione, sumoilazione

Silenziamento genico (lncRNAs, miRNAs)

Inattivazione del Cromosoma X

Imprinting genomico

Sistema Socio Sanitario

Cromosoma X

5% del DNA totale
155 milioni di paia di basi
800 geni codificanti
600 geni non codificanti (10% dei miRNA genomici totali)

Cromosoma Y

2% del DNA totale
58 milioni di paia di basi
75 geni codificanti
125 geni non codificanti

SRY: fattore di trascrizione che favorisce sviluppo dei testicoli. Regolatore espressione enzimi coinvolti metabolismo DA

Cromosoma X e risposta immunitaria

IMPRINTING GENOMICO

Table 1. Sex ratio between females and males in some neurodevelopmental and behavioural disorders.

Disease	Sex ratio [F:M]
Autism Spectrum Disorders (ASD)	1:3
Attention Deficit Hyperactivity Disorder (ADHD)	1:3
Schizophrenia	4:1
Dyslexia	1:1.6
Depression	1.7:1
Anxiety	1.7:1
Parkinson's disease (PD)	1:2
Alzheimer's disease (AD)	1.7:1

Saha et al, 2005⁴⁹; Miller et al, 2003⁵⁰; Albert 2015⁵¹; McLean et al, 2011⁵²; Loomes et al, 2017⁵³; Schmidt et al, 2008⁵⁴.

L'imprinting genomico consiste nell'espressione monoallelica (paterna o materna) di geni localizzati in regioni specifiche che vengono metilate

L'imprinting genomico avviene subito dopo la fecondazione ed ha un ruolo fondamentale nella crescita del feto e nello sviluppo della placenta

La perdita dell'imprinting causa patologie dello sviluppo fetale e disordini del comportamento (i.e. Angelman, Prader-Willi)

Il cervello contiene molte aree soggette ad imprinting

La maggior parte di patologie Mendeliane causate da mutazioni di proteine coinvolte nei processi epigenetici sono di natura neurologica (disabilità intellettiva, demenza)

Sistema Socio Sanitario

Differenze di genere nella genetica della Demenza di Alzheimer

Acta Neuropathol. 2018 Dec;136(6):857-872. doi: 10.1007/s00401-018-1881-4. Epub 2018 Jul 2.

Sex-specific genetic predictors of Alzheimer's disease biomarkers.

Deming Y¹, Dumitrescu L², Barnes LL³, Thambisetty M⁴, Kunkle B⁵, Gifford KA², Bush WS⁵, Chibnik LB^{6,7}, Mukherjee S⁸, De Jager PL^{9,10}, Kukull W¹¹, Huentelman M¹², Crane PK⁸, Resnick SM⁴, Keene CD¹³, Montine TJ¹⁴, Schellenberg GD¹⁵, Haines JL⁵, Zetterberg H^{16,17,18,19}, Blennow K^{16,17}, Larson EB^{8,20}, Johnson SC^{21,22}, Albert M²³, Moghekar A²³, Del Aguila JL¹, Fernandez MV¹, Budde J¹, Hassenstab J¹, Fagan AM²⁴, Riemenschneider M²⁵, Petersen RC²⁶, Minthon L²⁷, Chao MJ²⁸, Van Deerlin VM²⁹, Lee VM²⁹, Shaw LM²⁹, Trojanowski JQ²⁹, Peskind ER³⁰, Li G^{30,31}, Davis LK³², Sealock JM³², Cox NJ³²; Alzheimer's Disease Neuroimaging Initiative (ADNI); Alzheimer Disease Genetics Consortium (ADGC), Goate AM²⁸, Bennett DA³, Schneider JA³, Jefferson AL², Cruchaga C¹, Hohman TJ³³.

+ Collaborators (195)

+ Author information

Abstract

Cerebrospinal fluid (CSF) levels of amyloid- β 42 (A β 42) and tau have been evaluated as endophenotypes in Alzheimer's disease (AD) genetic studies. Although there are sex differences in AD risk, sex differences have not been evaluated in genetic studies of AD endophenotypes. ~~We performed sex-stratified and sex-interaction genetic analyses of CSF biomarkers to identify sex-specific associations. Data came from a previous genome-wide association study (GWAS) of CSF A β 42 and tau (1527 males, 1509 females).~~ We evaluated sex interactions at previous loci, ~~performed sex-stratified GWAS to identify sex-specific associations,~~ and evaluated sex interactions at sex-specific GWAS loci. We then evaluated sex-specific associations between prefrontal cortex (PFC) gene expression at relevant loci and autopsy measures of plaques and tangles using data from the Religious Orders Study and Rush Memory and Aging Project. In A β 42, we observed sex interactions at one previous and one novel locus: rs316341 within SERPINB1 ($p = 0.04$) and rs13115400 near LINC00290 ($p = 0.002$). These loci showed stronger associations among females ($\beta = -0.03$, $p = 4.25 \times 10^{-8}$; $\beta = 0.03$, $p = 3.97 \times 10^{-8}$) than males ($\beta = -0.02$, $p = 0.009$; $\beta = 0.01$, $p = 0.20$). Higher levels of expression of SERPINB1, SERPINB6, and SERPINB9 in PFC was associated with higher levels of amyloidosis among females (corrected p values < 0.02) but not males ($p > 0.38$). In total tau, we observed a sex interaction at a previous locus, rs1393060 proximal to GMNC ($p = 0.004$), driven by a stronger association among females ($\beta = 0.05$, $p = 4.57 \times 10^{-10}$) compared to males ($\beta = 0.02$, $p = 0.03$). There was also a sex-specific association between rs1393060 and tangle density at autopsy ($p_{\text{female}} = 0.047$; $p_{\text{male}} = 0.96$), ~~and higher levels of expression of two genes within this locus were associated with lower tangle density among females (OSTN $p = 0.006$; CLDN16 $p = 0.002$) but not males ($p \geq 0.32$).~~ Results suggest a female-specific role for SERPINB1 in amyloidosis and for OSTN and CLDN16 in tau pathology. Sex-specific genetic analyses may improve understanding of AD's genetic architecture.

Sistema Socio Sanitario

Differenze di genere nella genetica della Demenza di Alzheimer

JAMA Neurol. 2018 Aug 1;75(8):989-998. doi: 10.1001/jamaneurol.2018.0821.

Sex-Specific Association of Apolipoprotein E With Cerebrospinal Fluid Levels of Tau.

Hohman TJ¹, Dumitrescu L¹, Barnes LL², Thambisetty M³, Beecham G^{4,5}, Kunkle B⁵, Gifford KA¹, Bush WS⁶, Chibnik LB^{7,8}, Mukherjee S⁹, De Jager PL^{10,11}, Kukull W¹², Crane PK⁹, Resnick SM³, Keene CD¹³, Montine TJ¹⁴, Schellenberg GD¹⁵, Haines JL⁷, Zetterberg H^{16,17,18}, Blennow K^{16,17,19}, Larson EB^{9,20}, Johnson SC²¹, Albert M²², Bennett DA², Schneider JA², Jefferson AL¹; Alzheimer's Disease Genetics Consortium and the Alzheimer's Disease Neuroimaging Initiative.

CONCLUSIONS AND RELEVANCE: We provide robust evidence of a stronger association between APOE- ϵ 4 and CSF tau levels among women compared with men across multiple independent data sets. Interestingly, APOE- ϵ 4 is not differentially associated with autopsy measures of neurofibrillary tangles. ~~Together, the sex difference in the association between APOE and CSF measures of tau and the lack of a sex difference in the association with neurofibrillary tangles at autopsy suggest that APOE may modulate risk for neurodegeneration in a sex-specific manner, particularly in the presence of amyloidosis.~~

Sistema Socio Sanitario

Fondazione I.R.C.C.S.
Istituto Neurologico Carlo Besta

Regione
Lombardia

Differenze di genere nella genetica del Parkinson

Frontiers in Neuroendocrinology 50 (2018) 18–30

Contents lists available at ScienceDirect

Frontiers in Neuroendocrinology

journal homepage: www.elsevier.com/locate/yfrne

Review article

Sex differences in Parkinson's disease: Features on clinical symptoms, treatment outcome, sexual hormones and genetics

Juan Camilo Jurado-Coronel^a, Ricardo Cabezas^a, Marco Fidel Ávila Rodríguez^b,
Valentina Echeverría^{c,d}, Luis Miguel García-Segura^{c,f}, George E. Barreto^{a,g,*}

^a Departamento de Nutrición y Biotecnología, Facultad de Ciencias, Pontificia Universidad Javeriana, Bogotá D.C., Colombia

^b Facultad de Ciencias de la Salud, Universidad del Tolima, Ibagué, Colombia

^c Universidad San Sebastián, Fac. Cs de la Salud, Lientur 1457, Concepción, 4080871, Chile

^d Research & Development Service, Bay Pines VA Healthcare System, Bay Pines, FL 33744, USA

^e Instituto Cajal, CSIC, Madrid, Spain

^f CIBER de Investigación Biomédica en Red de Fragilidad y Envejecimiento Saludable (CIBERFES), Instituto de Salud Carlos III, Madrid, Spain

^g Instituto de Ciencias Biomédicas, Universidad Autónoma de Chile, Santiago, Chile

Parkinsonism and Related Disorders 20 (2014) 911–914

Contents lists available at ScienceDirect

Parkinsonism and Related Disorders

journal homepage: www.elsevier.com/locate/parkreldis

Short communication

LRRK2 mutations in Parkinson's disease: Confirmation of a gender effect in the Italian population

Roberto Cilia^a, Chiara Siri^a, Damiana Rusconi^b, Roberta Allegra^a, Andrea Ghiglietti^c,
Giorgio Sacilotto^a, Michela Zini^a, Anna L. Zecchinelli^a, Rosanna Asselta^d, Stefano Duga^d,
Anna M. Paganoni^c, Gianni Pezzoli^a, Manuela Seia^b, Stefano Goldwurm^{a,*}

^a Parkinson Institute, Istituti Clinici di Perfezionamento, via Bignami 1, 20126 Milan, Italy

^b Medical Genetics Laboratory, Foundation IRCCS "Ospedale Maggiore Policlinico, Mangiagalli e Regina Elena", Milan, Italy

^c MOX, Dipartimento di Matematica, Politecnico di Milano, Italy

^d Dipartimento di Biotechnologie Mediche e Medicina Traslazionale, Università degli Studi di Milano, Italy

Cervello:

SNCA e PINK1 up-regolati nell'uomo.

Geni coinvolti maturazione neuronale up-regolati nella donna.

Sistema Socio Sanitario

Fondazione I.R.C.C.S.
Istituto Neurologico Carlo Besta

Regione
Lombardia

THE LANCET

Volume 393 · Number 10171 · Pages 493-510 · February 9-15, 2019

www.thelancet.com

Advancing women in science, medicine, and global health

Editorial

Feminism is for everybody
See page 493

Articles

Are gender gaps due to evaluations of the applicant or the science?
See page 511

Articles

Sex-related reporting in medical research
See page 509

Review

Organisational best practices towards gender equality in science and medicine
See page 587

Health Policy

More talk than action: gender and ethnic diversity in universities
See page 594

£5.00 Registered as a newspaper · ISSN 0140-6736
Founded 1823 · Published weekly

Review

Gender equality in science, medicine, and global health: where are we at and why does it matter?

Geordan Shannon, Melanie Jansen, Kate Williams, Carlos Cáceres, Angelica Motta, Aloyce Odhiambo, Alie Ebeveld, Jenevieve Mannell

Lancet 2019; 393: 560-69 The purpose of this Review is to provide evidence for why gender equality in science, medicine, and global health

Correspondence

Addressing women's under-representation in medical leadership

self-belief of their male counterparts. Overall, for physicians seeking national awards, women account for 32% of applicants for bronze awards, 24% for

leaders. This finding, combined with the outpouring of positive feedback about the exhibition⁴ led to the commissioning of an RCP fellowship

73 | Special article

Ital J Gender-Specific Med 2018; 4(2): 73-78

Gender inequalities in medical careers: findings from five hospitals in the Lombardy Region

Camilla Gaiaschi

Department of Social and Political Sciences, University of Milan. Received 7 June 2018; accepted 7 September 2018.

Sistema Socio Sanitario

Fondazione I.R.C.C.S.
Istituto Neurologico Carlo Besta

Regione
Lombardia

**GUIDA AI CAMBIAMENTI STRUTTURALI
NEL MONDO ACCADEMICO
E NELLE ORGANIZZAZIONI DI RICERCA
PASSO DOPO PASSO**

| Regioni

L'importanza del Gender equality plan negli enti di cura e di ricerca

FRRB affronta l'argomento della parità di genere nell'ambito del progetto Target con la sua Community of practice. *IN COLLABORAZIONE CON FRRB

di [Redazione Aboutpharma Online](#)

21 marzo 2019

Sistema Socio Sanitario

Fondazione I.R.C.C.S.
Istituto Neurologico Carlo Besta

Regione
Lombardia

Con il termine “parità di genere” non si intende che uomini e donne debbano essere uguali o in ugual numero, ma che abbiano pari accesso alle stesse opportunità, pur conservando le loro peculiarità

CONSIGLIO DI AMMINISTRAZIONE

DELIBERAZIONE N. **IV/256**

Seduta n. ~~35~~ ³⁶ del 5 Dicembre 2018
~~35~~ del 12 Dicembre 2018

Presiede il Presidente

Alberto Guglielmo

Consiglieri

Andrea Casini
Ivano Loatelli
Adriano Paroli
Francesco Bevere
Agostino Migone De Amicis
Paolo Lazzati

Con l'assistenza del Segretario:

Paolo Tafuro

Su proposta : Direttore Generale

Germano Pellegata

Oggetto: Approvazione proposta Gender Equality Plan (Piano per l'Uguaglianza di Genere). Senza Oneri per la Fondazione.

Il Direttore Scientifico
Fabrizio Tagliavini

Il Direttore Amministrativo
Maurizia Ficarelli

Il Direttore Sanitario
Angelo Cordone

Il Responsabile del procedimento: Il Direttore U.O.C. Risorse Umane - Marco ~~Scos~~

Visto: Il Direttore del Dipartimento Amministrativo - Marco ~~Scos~~

L'atto si compone di n. 24 pagine, di cui n. 17 pagine di allegati, parte integrante

Grazie per l'attenzione

“What's the matter?
It's the same distance!”

Sistema Socio Sanitario

Fondazione I.R.C.C.S.
Istituto Neurologico Carlo Besta

Regione
Lombardia